

## ECUACIONES DE GRADO SUPERIOR A 2

**Resolver las siguientes ecuaciones:**

1)  $x^3 + 3x^2 - x - 3 = 0$

$$\begin{array}{c|cccc} & 1 & 3 & -1 & -3 \\ \hline 1 & & 1 & 4 & 3 \\ \hline & 1 & 4 & 3 & 0 \\ -1 & & -1 & -3 & \\ \hline & 1 & 3 & 0 & \end{array} \quad x^3 + 3x^2 - x - 3 = (x - 1)(x + 1)(x + 3)$$

$$(x - 1)(x + 1)(x + 3) = 0 \Rightarrow \begin{cases} x - 1 = 0 \Rightarrow x_1 = 1 \\ x + 1 = 0 \Rightarrow x_2 = -1 \\ x + 3 = 0 \Rightarrow x_3 = -3 \end{cases}$$

2)  $x^3 + 2x^2 + 2x + 1 = 0$

$$\begin{array}{c|cccc} & 1 & 2 & 2 & 1 \\ \hline -1 & & -1 & -1 & -1 \\ \hline & 1 & 1 & 1 & 0 \end{array} \quad x^3 + 2x^2 + 2x + 1 = (x + 1) \cdot (x^2 + x + 1)$$

$$(x - 1)(x^2 + x + 1) = 0 \Rightarrow \begin{cases} x - 1 = 0 \Rightarrow x_1 = 1 \\ x^2 + x + 1 = 0 \end{cases}$$

Las otras dos raíces las calculamos aplicando la fórmula de la ecuación de segundo grado:

$$x^2 + x + 1 = 0 \Rightarrow x = \frac{-1 \pm \sqrt{1-4}}{2} \rightarrow \text{No tiene solución}$$

3)  $x^3 + 3x^2 - 4x - 12 = 0$

$$\begin{array}{c|cccc} & 1 & 3 & -4 & -12 \\ \hline 2 & & 2 & 10 & 12 \\ \hline & 1 & 5 & 6 & 0 \\ -2 & & -2 & -6 & \\ \hline & 1 & 3 & 0 & \end{array} \quad x^3 + 3x^2 - 4x - 12 = (x - 2)(x + 2)(x + 3)$$

$$(x - 2)(x + 2)(x + 3) = 0 \Rightarrow \begin{cases} x - 2 = 0 \Rightarrow x_1 = 2 \\ x + 2 = 0 \Rightarrow x_2 = -2 \\ x + 3 = 0 \Rightarrow x_3 = -3 \end{cases}$$

4)  $x^3 - x^2 - x + 1 = 0$

$$\begin{array}{c|cccc} & 1 & -1 & -1 & 1 \\ \hline -1 & & -1 & 2 & -1 \\ \hline & 1 & -2 & 1 & 0 \\ 1 & & 1 & -1 & \\ \hline & 1 & -1 & 0 & \end{array} \quad x^3 - x^2 - x + 1 = (x + 1)(x - 1)^2$$

$$(x + 1)(x - 1)^2 = 0 \Rightarrow \begin{cases} x - 1 = 0 \Rightarrow x_1 = 1 \\ x + 1 = 0 \Rightarrow x_2 = -1 \end{cases}$$

5)  $x^3 - 2x^2 - 4x + 8 = 0$

$$\begin{array}{c|cccc} & 1 & -2 & -4 & 8 \\ \hline 2 & & 2 & 0 & -8 \\ \hline & 1 & 0 & -4 & 0 \\ 2 & & 2 & 4 & \\ \hline & 1 & 2 & 0 & \end{array} \quad x^3 - 2x^2 - 4x + 8 = (x + 2)(x - 2)^2$$

$$(x + 2)(x - 2)^2 = 0 \Rightarrow \begin{cases} x - 2 = 0 \Rightarrow x_1 = 2 \\ x + 2 = 0 \Rightarrow x_2 = -2 \end{cases}$$

6)  $6x^3 + 7x^2 - 9x + 2 = 0$

$$\begin{array}{c|ccccc} & 6 & 7 & -9 & 2 \\ \hline -2 & & -12 & 10 & -2 \\ \hline & 6 & -5 & 1 & 0 \end{array} \quad 6x^3 + 7x^2 - 9x + 2 = (x+2)(6x^2 - 5x + 1)$$

$$(x+2)(6x^2 - 5x + 1) = 0 \Rightarrow \begin{cases} x+2=0 \Rightarrow x_1 = -2 \\ 6x^2 - 5x + 1 = 0 \end{cases}$$

$$6x^2 - 5x + 1 = 0 \Rightarrow x = \frac{5 \pm \sqrt{5^2 - 4 \cdot 6 \cdot 1}}{6 \cdot 2} = \frac{5 \pm 1}{12} = \begin{cases} x_2 = \frac{5+1}{12} = \frac{6}{12} \Rightarrow x_2 = \frac{1}{2} \\ x_3 = \frac{5-1}{12} = \frac{4}{12} \Rightarrow x_3 = \frac{1}{3} \end{cases}$$

7)  $x^4 - 1 = 0$

$$x^4 - 1 = 0 \Rightarrow x^4 = 1 \Rightarrow x = \sqrt[4]{1} = \pm \Rightarrow [x = \pm 1]$$

8)  $8x^3 - 14x^2 + 7x - 1 = 0$

$$\begin{array}{c|ccccc} & 8 & -14 & 7 & -1 \\ \hline 1 & & 8 & -6 & 1 \\ \hline & 8 & -6 & 1 & 0 \end{array} \quad 8x^3 - 14x^2 + 7x - 1 = (x+2)(8x^2 - 6x + 1)$$

$$(x-1)(8x^2 - 6x + 1) = 0 \Rightarrow \begin{cases} x-1=0 \Rightarrow x_1 = 1 \\ 8x^2 - 6x + 1 = 0 \end{cases}$$

$$8x^2 - 6x + 1 = 0 \Rightarrow x = \frac{6 \pm \sqrt{6^2 - 4 \cdot 8 \cdot 1}}{8 \cdot 2} = \frac{6 \pm 2}{16} = \begin{cases} x_2 = \frac{6+2}{16} = \frac{8}{16} \Rightarrow x_2 = \frac{1}{2} \\ x_3 = \frac{6-2}{16} = \frac{4}{16} \Rightarrow x_3 = \frac{1}{4} \end{cases}$$

9)  $2x^4 - 5x^3 + 5x - 2 = 0$

$$\begin{array}{c|ccccc} & 2 & -5 & 0 & 5 & -2 \\ \hline 1 & & 2 & -3 & -3 & 2 \\ \hline & 2 & -3 & -3 & 2 & 0 \\ \hline -1 & & -2 & 5 & -2 & \\ \hline & 2 & -5 & 2 & 0 & \end{array} \quad 2x^4 - 5x^3 + 5x - 2 = (x+1)(x-1)(2x^2 - 5x + 2)$$

$$(x+1)(x-1)(2x^2 - 5x + 2) = 0 \Rightarrow \begin{cases} x-1=0 \Rightarrow x_1 = 1 \\ x+1=0 \Rightarrow x_2 = -1 \\ 2x^2 - 5x + 2 = 0 \end{cases}$$

$$2x^2 - 5x + 2 = 0 \Rightarrow x = \frac{5 \pm \sqrt{5^2 - 4 \cdot 2 \cdot 2}}{2 \cdot 2} = \frac{5 \pm 3}{4} = \begin{cases} x_3 = \frac{5+3}{4} = \frac{8}{4} \Rightarrow x_3 = 2 \\ x_4 = \frac{5-3}{4} = \frac{2}{4} \Rightarrow x_4 = \frac{1}{2} \end{cases}$$

10)  $x^4 + 2x^2 + 3 = 0$

$$y^2 + 2y + 3 = 0 \Rightarrow y = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 3}}{2} \Rightarrow [\text{No hay solución real}]$$